

Non-coincident Inter-instrument Comparisons of Ozone Measurements Using Quasi-conservative Coordinates

L. R. Lait¹, P. A. Newman², M. R. Schoeberl², T. McGee², L. Twigg¹, E. Browell³,
R. Bevilacqua⁴, S. B. Andersen⁵, H. DeBacker⁵, A. Benesova⁶, P. Skrivankova⁶, J.
Davies⁷, H. Claude⁸, E. Kyrö⁹, R. Kivi⁹, M. Gil¹⁰, M. Gerding¹¹, L. Zenobia¹², D.
Moore¹³, G. Braathen¹⁴, V. Dorokhov¹⁶, and I. Zaitcev¹⁶

¹Science Systems and Applications, Inc.

²NASA Goddard Space Flight Center

³NASA Langley Research Center

⁴U. S. Naval Research Laboratory

⁵Danish Meteorological Institute

⁶Czech Hydrometeorological Institute

⁷Meteorological Service of Canada

⁸Deutscher Wetterdienst Meteorologisches Observatorium

⁹Finnish Meteorological Institute

¹⁰Instituto Nacional de Técnica Aeroespacial, Spain

¹¹Leibniz-Institute of Atmospheric Physics, Germany

¹²Institute of Meteorology and Water Management, Poland

¹³U. K. Meteorological Office

¹⁴Norsk Institutt for Luftforskning

¹⁶Central Aerological Observatory, Russia

Manuscript submitted to

Atmospheric Chemistry and Physics

Manuscript version from May 24, 2004

Abstract

Ozone measurements from ozonesondes, AROTAL, DIAL, and POAM III instruments during the SOLVE-2/VINTERSOL period are composited in a time-varying, flow-following quasi-conservative (PV- θ) coordinate space; the resulting composites from each instrument are mapped onto the other instruments' locations and times. The mapped data are then used to intercompare data from the different instruments. Overall, the four data sets are found to be in good agreement. AROTAL shows somewhat lower values below 16 km, and DIAL has a positive bias at the upper limits of its altitude range. These intercomparisons are consistent with those obtained from more conventional near-coincident profiles, where available. Although the PV- θ mapping technique entails larger uncertainties of individual profile differences compared to direct near-coincident comparisons, the ability to include much larger numbers of comparisons can make this technique advantageous.

Correspondence to: Leslie R. Lait (lait@code916.gsfc.nasa.gov)

1 Introduction

The Kiruna deployments of the SOLVE-2 and VINTERSOL field experiments took place in January and February 2003. During this period, a number of different instruments measured stratospheric ozone. On board the NASA DC-8 aircraft were the Airborne Raman Ozone, Temperature, and Aerosol Lidar (AROTAL) and the Differential Absorption Lidar (DIAL), as well as in situ instruments such as FASTOZ and the Gas and Aerosol Measurement System/Langley Airborne Measurement Spectrometer (GAMES/LAABS). Other instruments, such as the balloon-borne MkIV interferometer, flew on other platforms or took ground-based measurements. A number of sites launched ozonesondes in coordination with the campaigns. In addition, data from the Polar Ozone and Aerosol Measurement III (POAM III) solar extinction instrument on the SPOT-4 spacecraft were made available to the mission teams.

Flying on board the NASA DC-8 together, the AROTAL and DIAL instruments were coincident, and the DC-8 flew over Ny Ålesund in coordination with a number of the sonde launches there. But because many of the instruments operated on different platforms at different times and locations, opportunities for measurement intercomparison were less than plentiful for most instruments. Aside from the AROTAL/DIAL data, the small number of near-coincident measurement sets make a statistical evaluation of inter-instrument differences problematic.

Other approaches are possible that can compare non-coincident data. The trajectory-mapping approach of Morris et al. (2000), the trajectory-hunting method of Danilin et al. (2003), and the MATCH technique of Rex et al. (1999) are examples.

In this work, a quasi-conservative coordinate method is employed. Described in Schoeberl and Lait (1991), this technique depends upon the premise that a reasonably long-lived trace gas should be well-mixed along contours of potential vorticity (PV) on a surface of constant potential temperature (θ) (Leovy et al. , 1985). By using PV and θ as coordinates, averaging mixing ratios near a set of points in that coordinate space should yield an accurate picture of a time-invariant trace gas distribution in PV- θ space, in the absence of diabatic effects and chemical changes. For short time periods, these effects often may safely be ignored; for periods longer than 10 days or so, they must be taken into account somehow.

PV- θ analysis was used in Schoeberl et al. (1989) and Lait et al. (1990) to map measurements onto a three-dimensional field, and in Kyrö et al. (2000) and Lait et al. (2002) to

determine stratospheric ozone loss in the Arctic. The quasi-conservative coordinate method can be also useful in inter-instrument comparisons, as seen in Redaelli et al. (1994), Manney et al. (2001), and Randall et al. (2002).

In this work, we intercompare ozone measurements from four instruments: AROTAL, DIAL, POAM III, and the ozonesondes. Section 2 describes each of these data sets briefly; then the analysis technique is described in section 3. Results follow in section 4.

2 Data

The SOLVE-2/VINTERSOL joint field experiment took place in January through early February 2003. We used the data from January 1 through February 10.

The NASA Goddard Space Flight Center's AROTAL instrument is a lidar that uses Rayleigh scattering from Eximer and Yag lasers transmitting at 308 and 355 nm to measure ozone, temperature, and aerosols. A more complete discussion of the instrument may be found in McGee et al. (2001) and Burris et al. (2002). For the SOLVE-2/VINTERSOL mission archive, AROTAL reports profiles every 22 seconds, averaged over 1.2 minutes. The altitude of the profiles depends on the altitude of the DC-8 aircraft, but over the middle of the flight it tends to range from approximately 14 km to 35 km. Vertical resolution of the reported data is approximately 150 m. Data were collected for 12 flights of the DC-8. To avoid problems with sunlight increasing noise in the measurements, only profiles taken where the local solar zenith angle is greater than 95° were used.

A second lidar, the NASA Langley Research Center's DIAL instrument, also flew on the DC-8. This instrument uses two YAG lasers transmitting at multiple frequencies to observe ozone below and above the aircraft. The DIAL data in the mission archive consist of profiles spaced about a minute apart. As with AROTAL, the altitudes covered change with the altitude of the aircraft, but typical coverage is from a few kilometers above the surface to around 25 km, with a small altitude gap near the aircraft itself. Vertical resolution reported is approximately 75 m. Data were collected for 14 flights (including two pre-mission test flights before the deployment to Kiruna). Details of this instrument may be found in Browell et al. (2003), Browell et al. (1998), and Richter et al. (1997).

A total of 213 balloon-launched sonde profiles from 21 ground stations were used in this analysis. These included special sondes launched for VINTERSOL, as well as those launched by the Meteorological Service of Canada, the World Meteorological Organization

network, Japan, and Russia. Data were used from the stations at Alert, Churchill, Eureka, Goosebay, Resolute, Stonyplain, Hohenpeissenberg, Jokioinen, Sodankyla, Keflavik, Kuehlungsborn, Legionowo, Lerwick, Ny Ålesund, Orland, Prague, Scoresbysund, Thule, Uccle, Salekhard, and Yakutsk. Altitude ranges vary greatly, but the sondes got as high as 29 km. Reported vertical resolution also varies, from about 10 m to around 60 m.

The POAM III solar occultation instrument is described in Lucke et al. (1999). It is a nine-channel photometer that uses solar occultation to measure atmospheric extinction in bands from 0.354 to 1.018 μm to retrieve temperature and multiple species, including ozone. Fourteen profiles are taken each day, spaced around a latitude circle that moves slowly in time. Vertical resolution is approximately 1 km (Lumpe et al., 2003).

To apply the quasi-conservative coordinate analysis, values of potential vorticity and potential temperature must be obtained at each measurement location and time. For this work, these are obtained by interpolating three-dimensional gridded analyses from the Data Assimilation Office (now the Global Modelling and Assimilation Office) of NASA's Goddard Space Flight Center. These analyses were chosen because of their relatively fine horizontal and temporal resolution, as well as their altitude range. The analyses used were chosen from the "final look" product generated from the GEOS-4 system. GEOS-4 was the successor to the GEOS-1 system documented by Pfaendner et al. (1995). These data grids extend from 1000 to 0.2 hPa, have a horizontal resolution of 1.25° longitude by 1.0° latitude, and are produced four times daily.

3 Analysis

In order to derive meaningful statistics for both coincident and non-coincident comparisons, it was necessary to obtain uncertainties associated with the ozone measurements. To estimate these uncertainties for AROTAL and DIAL, a standard variance profile for each instrument was constructed from the small-scale horizontal variations of the all profiles from all flights. For ozonesondes, we used the larger of 5% of the measured value, or the variance about a linear fit within a 5 km segment. Uncertainties in the POAM III measurements were supplied in the data archive.

The measurements, with their uncertainties, locations, and times were collected for each instrument. (Note that for the sondes, the winds were used to estimate the horizontal location of each balloon during its ascent.) Modified potential vorticity (see Lait (1994)) and

potential temperature were interpolated from the meteorological analysis onto the measurement locations and times.

By computing autocorrelations of the lidar data along the flight, we were able to estimate the minimum horizontal separation to ensure independence of profiles as approximately 400 km for AROTAL measurements and 375 km for DIAL. These distances are roughly consistent with those used by Schoeberl et al. (2002). Because the profile sites for the sondes and POAM III data were widely separated, horizontal separations were not an issue for those instruments.

Minimum vertical separations between a profile's measurements were similarly obtained. For AROTAL, the vertical separation was estimated to be 3 km; for DIAL, it was 4 km. For the sondes, it was 3 km, and for POAM data, 4 km.

Measurements closer than these horizontal and vertical separations were omitted from the analysis, so that the data going into the analysis could be considered independent of each other.

The analysis itself is similar to that in Lait et al. (2002). A regular grid in a PV- θ coordinate space was constructed, and PV and θ values interpolated from the analyses were used to locate each ozone measurement in the coordinate space. Data from January 1 through February 10 were used; hence, diabatic effects and chemical changes needed to be accounted for. These effects both show up as a change in ozone over time at a given point in PV- θ space. To first order, they can be dealt with by applying a weighted linear time fit to the data near a given PV- θ gridpoint. Each point was weighted inversely to its uncertainty and its distance from the PV- θ gridpoint being examined.

This procedure yields a set of slopes and intercepts, one for each gridpoint in the PV- θ coordinate space. From these, we can construct a composite field in PV- θ space for any moment in time; moreover, given the meteorological analyses we can map this ozone field back into longitude-latitude-altitude space,.

Note that for instrument comparison purposes here, only the overall evolution of the ozone field is relevant. Whether changes in that field are caused by diabatic effects or by chemical loss/production is of no concern, and so there is no need to try to separate those two effects.

Data from two instruments are compared by constructing a PV- θ composite field from one instrument and mapping its ozone values onto the locations and times of the second instrument. The differences between the two ozone values and the uncertainties associated

with those differences are collected, and mean profiles of the biases are computed for the mission period, taking the uncertainties into account.

4 Results

To validate the analysis technique, several tests were applied.

First, each instrument was compared against itself. That is, differences were characterized between the original measurements and those from PV- θ mapping of the same instrument's data. This test should reveal any biases or distortions introduced by the analysis technique itself, and it should also reveal the degree to which noise is introduced by errors in the meteorological fields, departures from the well-mixedness assumption, and so on. Figure 1 shows an example of the self-comparison for AROTAL data. The maximum difference, 0.1 ppmv near 22 km, is not statistically different from zero, and the rest of the profile is very close to zero. The self-comparisons for the other instruments show similar results: very small average differences, with at most minor statistically insignificant fluctuations.

The next test was to compute inter-instrument differences between AROTAL and DIAL using both near-coincident and non-coincident methods. Because both these instruments flew aboard the same aircraft, a large number of near-coincident profiles could be collected. For the 12 SOLVE-2 flights of the DC-8, the two closest profiles of the two instruments were chosen; each profile pair also had to be separated from all other profile pairs by at least 400 km. For each profile pair, the DIAL data were then interpolated to the AROTAL altitudes, and the two profiles were differenced.

Figure 2 shows the differences and their average profile. Above 20 km, DIAL ozone values are systematically higher than AROTAL. Below 16 km, AROTAL values are higher.

Comparing the AROTAL and DIAL data using the non-coincident PV- θ analysis yields similar results (Figure 3). Note that the measured-reconstructed differences are consistent with the reconstructed-measured differences. Of course, because the data being compared were in fact coincident, this is no more rigorous a test of the noncoincident technique than the self-comparisons. evaluating the success of the next test.

A more demanding test is to compare true non-coincident DIAL and AROTAL data. To accomplish this, AROTAL data from the even-numbered flights were compared with DIAL data from the odd-numbered flights, and vice-versa. The results, shown in Figures 4 and 5,

are consistent with the near-coincident comparisons, albeit with larger uncertainties (since they involved only half the data). Figure 4 matches the full data comparison quite well, while Figure 5 has greater uncertainties but is still roughly consistent with the others.

Having confirmed that each instrument's data compare well with themselves, and that the AROTAL-DIAL non-coincident comparisons are similar to the coincident comparisons, we proceeded to compare the other instruments' data.

Figure 6 shows the differences between AROTAL and the ozonesondes. There appears to be a slight bias below 15 km, with AROTAL being perhaps 0.3 to 0.4 ppmv higher than the sondes near 12–13 km. Above 25 km, the ozonesondes are fewer in number and their uncertainties are often larger, so that the error bars in the differences are much larger at those altitudes.

Likewise, the differences between DIAL and the sondes is shown in Figure 7. Here, DIAL matches the sondes well at the lower altitudes, but DIAL is higher at the uppermost reaches of the instrument, above 25 km. Note also that the AROTAL-sonde and DIAL-sonde differences are consistent with the AROTAL-DIAL differences.

Figures 8 and 9 show the POAM-AROTAL and POAM-DIAL differences, respectively. The POAM-AROTAL are qualitatively similar to biases noted in Lumpe et al (2003), where coincident comparisons were made between POAM III and AROTEL (an earlier version of AROTAL) and DIAL during the first SOLVE campaign in the winter of 1999-2000. The POAM-DIAL differences at the uppermost DIAL altitudes, however, are of different sign from those in Lumpe et al (2003)

Figure 10 compares the ozonesonde data with the POAM III profiles. A small bias appears near 20 km. At the uppermost altitudes, where the sonde measurements are fewer and less certain, there is also a suggestion of a small negative bias.

These intercomparisons were repeated with meteorological analysis products from other institutions, and the results were similar.

5 Conclusions

Ozone measurements taken during the SOLVE-2/VINTERSOL field experiment from four instruments (AROTAL, DIAL, POAM III, and sondes) were compared. A quasi-conservative coordinate approach was employed to use non-coincident data for instrument intercomparisons. Several tests of the method were applied. First, each instrument's data were self-

compared; the differences were zero, within the uncertainty associated with the technique. Second, the AROTAL and DIAL data were compared, with results similar to those from the straightforward near-coincident comparison. These tests demonstrate that the quasi-conservative technique does not introduce unusual or misleading artifacts into the data. To provide a true non-coincident data comparison, the DIAL data from the odd-numbered DC-8 flights were compared with AROTAL data from the even-numbered flights, and vice versa. These results were consistent with the other DIAL-AROTAL comparisons, although the uncertainties were larger.

Finally, all four data sets were intercompared. The AROTAL ozone values were found to be 0.1 to 0.4 ppmv higher than DIAL and the sondes below 16 km, and DIAL values were 0.3 to 0.8 ppmv higher than AROTAL and the sondes above 22 km. The ozonesonde data match well with POAM III, except for a possible small bias in a region near 22 km.

This PV- θ analysis produces results with substantial uncertainties. The uncertainties associated with the analyzed PV and θ meteorological fields, limited sampling over regions of PV- θ space, potential failures in the assumptions necessary for the method's validity (e.g., homogeneity around a circumpolar PV- θ tube), and even a breakdown in PV-ozone correlations at high altitudes and low latitude, can all contribute to the enlarged error bars. Certainly, then, direct comparison of large numbers of near-coincident measurements is preferred where it is possible. Nevertheless, the higher numbers of comparisons which are made possible by relaxing the requirement for near-coincidence, can improve the statistics so that the results are useful despite their uncertainties.

Acknowledgements.

We wish to acknowledge the DC-8 pilots and the flight and ground crews who made the aircraft measurements possible under sometimes difficult conditions; the sonde launch personnel; the SOLVE0-2 logistics staff and the personnel of the Arena Arctica in Kiruna, Sweden; and SOLVE-2 and VINTERSOL management. This research was supported by the NASA's Atmospheric Chemistry Modeling and Analysis Program and the Upper Atmosphere Research Program.

References

- Browell, E. V., S. Ismail, and W. B. Grant, Differential Absorption Lidar (DIAL) measurements from air and space, *Appl. Phys.-B*, 67, 399–410, 1998.
- Browell, E. V., M. A. Fenn, C. F. Butler, W.B. Grant, V. G. Brackett, J. W. Hair, M. A. Avery, R. E. Newell, Y. Hu, H. E. Fuelberg, D. J. Jacob, B. E. Anderson, E. L. Atlas, D. R. Blake, W. H. Brune, J. E. Dibb, A. Fried, B. G. Heikes, G. W. Sachse, S. T. Sandholm, H. B. Singh, R. W. Talbot, S. A. Vay, R. J. Weber, and K. B. Bartlett, Large-scale ozone and aerosol distributions, air mass characteristics, and ozone fluxes over the western Pacific Ocean in late winter/early spring, *J. Geophys. Res.*, 108, doi:10.1029/2002JD003290, 2003.
- Burris, J., T. McGee, W. Hoegy, L. Lait, L. Twigg, G. Sumnicht, W. Heaps, C. Hostetler, T. P. Bui, R. Neuber, I. S. McDerimid, Validation of temperature measurements from the airborne Raman ozone temperature and aerosol lidar during SOLVE, 107, doi:10.1029/2001JD001028, 2002.
- Danilin, M. Y., M. K. W. Ko, R. M. Bevilacqua, L. V. Lyjak, L. Froidevaux, M. L. Santee, J. M. Zawadny, K. W. Hoppel, E. C. Richard, J. R. Spackman, E. M. Weinstock, R. L. Herman, K. A. McKenney, P. O. Wennberg, F. L. Eisle, R. M. Stempfle, C. J. Scott, J. W. Elkins, and T. V. Bui, Comparison of ER-2 aircraft and POAM III, MLS, and SAGE II satellite measurements during SOLVE using traditional correlative analysis and trajectory hunting technique, *J. Geophysical Res.*, 108, 8315, doi:10.1029/2001JD000781, 2003.
- Kyrö, E., R. Kivi, T. Turunen, H. Aulamo, V. V. Rudakov, V. Khattatov, A. R. MacKenzie, M. P. Chipperfield, A. M. Lee, L. Stefanutti, and F. Ravegnani, Ozone measurements during the Airborne Polar Experiment: aircraft instrument validation, isentropic trends, and hemispheric fields prior to the 1997 Arctic ozone depletion, *J. Geophys. Res.*, 105, 14599–14611, 2000.
- Lait, L. R., M. R. Schoeberl, P. A. Newman, M. H. Proffitt, M. Lowenstein, J. R. Podolske, S. E. Strahan, K. R. Chan, B. Gary, J. J. Margitan, E. Browell, M. P. McCormick, and A. Torres, Reconstruction of O₃ and N₂O fields from ER-2, DC-8, and balloon observations, *Geophys. Res. Lett.*, 17, 521–524, 1990.
- Lait, L., An alternative form for potential vorticity, *J. Atmos. Sci.*, 51, 1754–1759, 1994.
- Lait, L. R., M. R. Schoeberl, P. A. Newman, T. McGee, J. Burris, E. V. Browell, E. Richard, G. O. Braathen, B. R. Bojkov, F. Goutail, P. von der Gathen, E. Kyrö, G. Vaughan, H. Kelder, S. Kirkwood, P. Woods, V. Dorokhov, I. Zaitcev, Z. Litynska, B. Kois, A. Benesova, P. Skrivankova, H. De Backer, J. Davies, T. Jorgensen, and I. S. Mikkelsen, Ozone loss from quasi-conservative coordinate mapping during the 1999–2000 SOLVE/THESEO 2000 Campaigns, *J. Geophys. Res.*, 107, doi:10.1029/2001JD000998, 2002.
- Leovy C. B., C-R. Sun, M. H. Hitchman, E. E. Remsberg, J. M. Russell III, L. L. Gordley, J. C. Gille, and L. V. Lyjak, Transport of ozone in the middle stratosphere: evidence for wave breaking. *J. Atmos. Sci.*, 42, 230–244, 1985.

- The Polar Ozone and Aerosol Measurement (POAM) III instrument and early validation results, *J. Geophys. Res.*, 104, 18785–18799, 1999.
- Lumpe, J. L., M. Fromm, K. Hoppel, R. M. Bevilacqua, C. E. Randall, E. V. Browell, W. B. Grant, T. McGee, J. Burris, L. Twigg, E. C. Richard, G. C. Toon, J. J. Margitan, B. Sen, K. Pfeilsticker, H. Boesch, R. Fitzenberger, F. Goutail, J.-P. Pommereau, Comparison of POAM III ozone measurements with correlative aircraft and balloon data during SOLVE, *J. Geophys. Res.*, 108, doi:10.1029/2001JD000472, 2003.
- Manney, G. L., H. A. Michelsen, R. M. Bevilacqua, M. R. Gunson, F. W. Irion, N. J. Livesy, J. Oberheide, M. Riese, J. M. Russell III, G. C. Toon, and J. M. Zawodny, Comparison of satellite ozone observations in coincident air masses in early November 1994, *J. Geophys. Res.*, 106, 9923–9943, 2001.
- McGee, T. J., M. Gross, U. N. Singh, J. J. Butler, and P. Kimvilankani, An improved stratospheric ozone lidar, *Opt. Eng.*, 20, 955–958, 1995.
- Morris, G. A., J. F. Gleason, J. Ziemke, M. R. Schoeberl, Trajectory mapping: a tool for validation of trace gas observations, *J. Geophys. Res.*, 105, 17875–17894, 2000.
- Pfaendtner, J., S. Bloom, D. Lamich, M. Seabloom, M. Sienkiewicz, J. Stobie, and A. da Silva, Documentation of the Goddard Earth Observing System (GEOS) Data Assimilation System - Version 1, NASA Technical Memorandum 104606, Vol. 4, 1995.
- Randall, C. E., J. D. Lumpe, R. M. Bevilacqua, L. W. Hoppel, M. D. Fromm, R. J. Salawitch, W. H. Swartz, S. A. Lloyd, E. Kyrö, P. von der Gathen, H. Claude, J. Davies, H. DeBacker, H. Dier, M. J. Molyneux, and J. Sancho, Reconstruction of three-dimensional ozone fields using POAM III during SOLVE, *J. Geophys. Res.*, 107, doi:10.1029/2001JD000471, 2002.
- Redaelli, G., L. R. Lait, M. Schoeberl, P. A. Newman, G. Visconti, A. Daltorio, F. Masci, V. Rizzi, L. Froidevaux, J. W. Waters, A. J. Miller, UARS MLS O₃ soundings compared with lidar measurements using the conservative coordinates reconstruction technique, *Geophys. Res. Lett.*, 21, 1535–1538, 1994.
- Rex, M., P. von der Gathen, G. O. Braathen, N. R. P. Harris, E. Reimer, A. Beck, R. Alfier, R. Krüger-Carstensen, M. Chipperfield, H. De Backer, D. Balis, F. O’Conner, H. Dier, V. Dorokhov, H. Fast, A. Gamma, M. Gil, E. Kyrö, Z. Litynska, I. S. Mikkelsen, M. Molyneux, G. Murphy, S. J. Reid, M. Rummukainen, and C. Zerefos, Chemical ozone loss in the Arctic winter 1994/1995 as determined by the Match technique, *J. Atmos. Chem.*, 32, 35–59, 1999.
- Richter, D. A., E. V. Browell, C. F. Butler, and N. S. Higdon, Advanced airborne UV DIAL system for stratospheric and tropospheric ozone and aerosol measurements, in *Advances in Atmospheric Remote Sensing with Lidar*, edited by A. Ansmann et al., pp. 395–398, Springer Verlag, New York, 1997.
- Schoeberl, M. R., L. R. Lait, P. A. Newman, R. L. Martin, M. H. Proffitt, D. L. Hartmann, M. Lowenstein, J. Podolske, S. E. Strahan, J. Anderson, K. R. Chan, and B. Gary, Reconstruction of the

constituent distribution and trends in the Antarctic polar vortex from ER-2 flight observations, *J. Geophys. Res.*, *94*, 16815–16845, 1989.

Schoeberl M. R. and L. R. Lait, Conservative coordinate transformations for atmospheric measurements, in *The Use of EOS for Studies in Atmospheric Physics*, edited by J. C. Gille and G. Visconti, *Proc. Internat. School of Phys. "Enrico Fermi", CXV Course*, 419–430, 1991.

Schoeberl, M. R., P. A. Newman, L. R. Lait, T. J. McGee, J. F. Burris, E. V. Browell, W. B. Grant, E. C. Richard, P. von der Gathen, R. Bevilacqua, I. S. Mikkelsen, M. J. Molyneux, An Assessment of the Ozone Loss During the 1999-2000 SOLVE/THESEO 2000 Arctic Campaign, *J. Geophys. Res.*, *107*, doi:10.1029/2001JD000412, 2002.

Figure Captions

Fig. 1. Self-comparison biases for AROTAL data, expressed the difference between the original measurements and the measurements reconstructed through PV- θ mapping. Dots are the individual differences, the thin line is the mean difference profile, and the thick horizontal lines are the 90% confidence limits of the mean differences. The average measured ozone profile is shown in the right panel.

Fig. 2. Near-coincident differences between AROTAL and DIAL. Gray dots represent individual differences, the solid line shows the mean difference profiles, horizontal lines show the 90% confidence limits of the mean profile, and the dotted lines show the standard deviation of the differences.

Fig. 3. Differences between AROTAL and DIAL, using PV- θ analysis. As in Fig. 1, the left panel shows the differences, and the right panel shows the average ozone profile. The mean differences (with their 90% confidence limits) are shown for the comparisons using AROTAL and DIAL as the data which is mapped onto the other instrument's locations.

Fig. 4. As in Fig. 3, except that AROTAL data from the even-numbered DC-8 flights are compared with DIAL data from the odd-numbered flights.

Fig. 5. As in Fig. 3, except that AROTAL data from the odd-numbered DC-8 flights are compared with DIAL data from the even-numbered flights.

Fig. 6. As in Fig. 3, except that AROTAL data are compared with data from the ozonesondes.

Fig. 7. As in Fig. 3, except that DIAL data are compared with data from the ozonesondes.

Fig. 8. As in Fig. 3, except that AROTAL data are compared with data from POAM III.

Fig. 9. As in Fig. 3, except that DIAL data are compared with data from POAM III.

Fig. 10. As in Fig. 3, except that ozonesonde data are compared with data from POAM III.

Figures




Fig. 1. Self-comparison biases for AROTAL data, expressed the difference between the original measurements and the measurements reconstructed through PV- θ mapping. Dots are the individual differences, the thin line is the mean difference profile, and the thick horizontal lines are the 90% confidence limits of the mean differences. The average measured ozone profile is shown in the right panel.




Fig. 2. Near-coincident differences between AROTAL and DIAL. Gray dots represent individual differences, the solid line shows the mean difference profiles, horizontal lines show the 90% confidence limits of the mean profile, and the dotted lines show the standard deviation of the differences.




Fig. 3. Differences between AROTAL and DIAL, using PV- θ analysis. As in Fig. 1, the left panel shows the differences, and the right panel shows the average ozone profile. The mean differences (with their 90% confidence limits) are shown for the comparisons using AROTAL and DIAL as the data which is mapped onto the other instrument's locations.




Fig. 4. As in Fig. 3, except that AROTAL data from the even-numbered DC-8 flights are compared with DIAL data from the odd-numbered flights.




Fig. 5. As in Fig. 3, except that AROTAL data from the odd-numbered DC-8 flights are compared with DIAL data from the even-numbered flights.




Fig. 6. As in Fig. 3, except that AROTAL data are compared with data from the ozonesondes.




Fig. 9. As in Fig. 3, except that DIAL data are compared with data from POAM III.




Fig. 10. As in Fig. 3, except that ozonesonde data are compared with data from POAM III.




Fig. 7. As in Fig. 3, except that DIAL data are compared with data from the ozonesondes.




Fig. 8. As in Fig. 3, except that AROTAL data are compared with data from POAM III.

Popular Summary of
"Non-coincident Inter-instrument Comparisons of Ozone
Measurements Using Quasi-conservative Coordinates"
by Lait, et al.

This paper compares stratospheric ozone measurements from four instruments during the SOLVE-2/VINTERSOL field experiment in January-February 2003. The four instruments are: the NASA Goddard Space Flight Center AROTAL airborne lidar, the NASA Langley Research Center DIAL airborne lidar, the satellite-based POAM III instrument of the U. S. Naval Research Lab, and various ozonesondes launched during the period. Because the number of coincident measurements was small for most instruments, a special non-coincident technique was employed here that uses conserved meteorological quantities to map data from one instrument onto another instrument's measurement locations. We find that most instruments agree reasonably well over most of their altitude ranges. AROTAL has a somewhat positive bias below around 13 km, and DIAL has a stronger positive bias above 22 km.